

Towards consistent theory of geometric origin with massive spin-2 field

V.A. Rubakov

Institute for Nuclear Research
of the Russian Academy of Sciences, Moscow

with V. Nikiforova (Moscow), V.P. Nair (New York)
and S. Randjbar-Daemi (Trieste)

0811.3781;

work in progress, to appear soon

Motivaton and Introduction

- **Modification of gravity at large distances**
Challenge for theory, motivated in particular by accelerated expansion of the Universe.
 - May accelerated expansion be due to new gravity at cosmological scales rather than due to new form of energy?
 - **Difficult to construct consistent theory**
 - **Prototype theory: Fierz–Pauli theory of massive graviton**
Linearized about Minkowski background:
 - van Dam–Veltman–Zakharov discontinuity
 - non-linearity at large distances from a source \iff
Vainshtein phenomenon
(actually may cure vDVZ problem)
- Disaster in curved backgrounds: Boulware–Deser ghost**

Approaches:

- Scalar–tensor gravities, $f(R)$ -gravities

Boisseau, Esposito–Farese, Polarski, Starobinsky' 00
Gannouji, Polarski, Ranquet, Starobinsky' 06,
etc.

- Extra dimensions with brane worlds

Dvali, Gabadadze, Porrati' 01
de Rham et. al.' 07
Kaloper, Kiley' 07
Kobayashi' 07
etc.

- Lorentz-violating effective Lagrangians

V.R.'04
Dubovsky' 04
Dubovsky, Tinyakov, Tkachev' 05
Berezhiani, Comelli, Nesti, Pilo' 07
etc.

Some proposals work: apparently consistent theories with modified Newton's law at large distances.

NB: Obtaining accelerating Universe in a "natural" way is still difficult

Old work

Hayashi, Shirafuji' 80

Sezgin, van Nieuwenhuizen' 80

- Theories with both metric and connection dynamical fields.
 - Many have both massive and massless spin-2 fields at linearized level in Minkowski background
 - Tuning of parameters \implies no pathologies in Minkowski background (no ghosts, no tachyons)

Suspicious: in curved backgrounds tunings may be ruined \implies pathologies may reappear

This is precisely what happens in Fierz–Pauli gravity \implies Boulware–Deser ghost mode

This work: look into this issue using brute force approach

Outcome: No, theory remains healthy at least in Einstein backgrounds of weak enough curvature

Model

● Field content

- metric $g_{\mu\nu} \implies$ vierbein e_{μ}^i
- connection $A_{ij\mu} =$ gauge field of local $O(3,1)$, Lorentz group of frame rotations $\implies A_{ijk} \equiv e_i^{\mu} A_{ij\mu} = -A_{jik}$.

● Covariant quantities:

- Curvature = Field strength, just like in Yang–Mills:

$$F_{ij\mu\nu} = \partial_{\mu} A_{ij\nu} - \partial_{\nu} A_{ij\mu} + A_{ik\mu} A^k_{j\nu} - A_{ik\nu} A^k_{j\mu}$$

$$F_{ijmn} = e_m^{\mu} e_n^{\nu} F_{ij\mu\nu}, \quad \text{frame basis}$$

- Torsion \simeq (Connection – Riemann connection)

$$A_{ijk} = \frac{1}{2} (T_{ijk} - T_{jik} - T_{kij} + C_{ijk} - C_{jik} - C_{kij})$$

$$C_{ijk} = e_j^\mu e_k^\nu (\partial_\mu e_{i\nu} - \partial_\nu e_{i\mu}) = -C_{ikj} \iff \text{Riemann connection}$$

T_{ijk} = torsion tensor.

- Gauge symmetries:
 - Local Lorentz
 - General coordinate

- Trick to construct theories with massive fields in gauge-invariant way:

$$L = (F + F^2) + T^2$$

$$L = (F + F^2) + T^2$$

- One way to look:

$F = \partial_\mu A \implies F^2 =$ kinetic term for connection A_{ijk}

$$T = A - C = A - \partial_\mu e \implies$$

$T^2 =$ mass term for connection A_{ijk} plus kinetic term for vierbein e_μ^i (plus mixing between the two).

- Another way to look:

$$A = T + C = T + \partial_\mu e \implies$$

$F^2 =$ higher order kinetic term for vierbein e_μ^i plus kinetic term for torsion T_{ijk} (plus mixing)

$T^2 =$ mass term for torsion

Signs not guaranteed \implies
Danger of ghosts and tachyons

- Number of fields: 10 from e^i_μ + 24 from T_{ijk} = 34.
- Number of gauge symmetries: 4 of **general coordinate transformations** + 6 of **local Lorentz** = 10
- Generally: number of propagating degrees of freedom:

$$34 - 2 \times 10 = 14$$

Some of them **ghosts**

- Parameters fine tuned: less propagating modes,
no ghosts about Minkowski background

Hayashi, Shirafuji' 80
Sezgin, van Nieuwenhuizen' 80

In what follows: one of the cases

- Decomposition of torsion into irreps of $O(3, 1)$:

$$T_{ijk} = \frac{2}{3}(t_{ijk} - t_{ikj}) + \frac{1}{3}(\eta_{ij}v_k - \eta_{ik}v_j) + \varepsilon_{ijkl}a^l$$

with $t_{ijk} = t_{jik}$ and

$$t_{ijk} + t_{jki} + t_{kij} = 0, \quad \eta^{ij}t_{ijk} = 0, \quad \eta^{ik}t_{ijk} = 0$$

- **Fine tuning the parameters:** $L = L_{F+F^2} + L_{T^2}$

“Kinetic term”:

$$L_{F+F^2} = M^2 F + \lambda + c_3 F_{ij} F^{ij} + c_4 F_{ij} F^{ji} + c_5 F^2 + c_6 (\varepsilon_{ijkl} F^{ijkl})^2$$

$$(1): \quad c_3 + c_4 + 3c_5 = 0, \quad (2): \quad \text{no term } F_{ijkl} F^{ijkl}$$

“Mass term”:

$$(3): \quad L_{T^2} = -\mu^2 (t_{ijk} t^{ijk} - v_i v^i + \frac{9}{4} a_i a^i)$$

Parameters:

- M^2, μ^2 : dimension (mass)².
Assume for the sake of presentation $M \sim \mu \sim M_{Pl}$.
- c_3, c_4, c_5, c_6 : dimensionless.
Assume for the sake of presentation $c_3 \sim c_4 \sim c_5 \sim c_6 \equiv c$
- Signs:

$$c_5 < 0, \quad c_6 > 0, \quad 0 < \mu^2 < \frac{2}{3}M^2$$

This ensures absence of ghosts and tachyons in Minkowski background.

- λ : cosmological constant, introduced for generality: useful for discussing Einstein spaces as backgrounds, not just Ricci flat

Field equations

$$\frac{\delta S}{\delta e_{\mu}^i} : \quad M^2 F_{ij} + \{c_i\} (F \cdot F)_{ij} + (\text{torsion})_{ij} - (1/2)L = 0$$

$$\frac{\delta S}{\delta A_{ijk}} : \quad \{c_i\} (DF)_{ijk} + (\text{torsion})_{ijk} = 0$$

Involve F_{ij} , do not involve F_{ijkl} .

Consequences

- $\lambda = 0 \iff$ torsion = 0, $F_{ij} = R_{ij} = 0$ always a solution.
- Massless tensor field — graviton from e_{μ}^i — in Minkowski background.
- $\lambda \neq 0 \iff$ Einstein space with zero torsion, $R_{ij} = \Lambda g_{ij}$ also a solution

$$\Lambda = \frac{\lambda}{M^2} \quad \text{just like in GR.}$$

Spectrum about Minkowski background

Hayashi, Shirafuji' 80

Sezgin, van Nieuwenhuizen' 80

- Massless spin-2 from e_{μ}^i
- Vector v_i : not a propagating field
- Transverse part of pseudovector a_i : also does not propagate.
- Longitudinal part of a_i : massive pseudoscalar field,

$$m_1^2 \sim \frac{M^2}{c}$$

- Propagating part of t_{ijk} mixed with e_{μ}^i : massive spin-2 field

$$m^2 \sim \frac{M^2}{c}$$

NB: Number of propagating modes reduced from 14 to 8.

Sources in Minkowski background

- Notation for perturbation of vierbein:

$$e_{\mu}^i = \delta_{\mu}^i + h_{\mu}^i$$

- General source term

$$S_{source} = \int d^4x \left(2h_{\mu}^i \theta_i^{\mu} - \frac{1}{2} A_{ij\mu} S^{ij\mu} \right)$$

Gauge invariance \implies

$$\partial^j \theta_{ij} = 0 \implies \partial^j \theta_{(ij)} = -\partial^j \theta_{[ij]}$$

$$\partial_k S^{ijk} = 4\theta^{[ij]}$$

Fields induced by sources:

$$h_{ij} = \frac{1}{M^2} \frac{1}{k^2} \left(\tau_{ij} - \frac{1}{2} \eta_{ij} \tau \right) + \frac{2M^2/3 - \mu^2}{\mu^2 M^2} \frac{1}{k^2 + m^2} \left(\sigma_{ij} - \frac{1}{3} \eta_{ij} \sigma + \text{more terms with } \sigma \right)$$

with

$$\tau^{ij} = \theta^{(ij)} - \frac{1}{2} \partial_k S^{k(ij)}$$

$$\sigma^{ij} = \theta^{(ij)} + \frac{\mu^2}{2M^2/3 - \mu^2} \frac{1}{2} \partial_k S^{k(ij)}$$

- Sum of massless and massive spin-2 contributions
 - same for $S_{ijk} = 0$
- vDVZ discontinuity in massive spin-2 sector
- Mixing with torsion \iff Source for torsion contributes to metric

Curved backgrounds

Digression: Boulware–Deser mode in Fierz–Pauli theory

Fierz–Pauli action:

$$S = M_{\text{Pl}}^2 \int d^4x \sqrt{g} \left\{ L_{EH}(g_{\mu\nu}) - \frac{m^2}{4} h_{\mu\nu} h^{\mu\nu} + \frac{m^2}{4} (h^\mu{}_\mu)^2 + \# \cdot h^3 + \dots \right\}$$

with

$$h_{\mu\nu} = g_{\mu\nu} - \eta_{\mu\nu}$$

- Danger: explicitly broken gauge invariance
- How to extract dangerous degrees of freedom?

Stückelberg trick

Arkani-Hamed, Georgi, Schwartz' 03
Creminelli, Nicolis, Papucci, Trincherini' 05
Deffayet, Rombouts' 05

- L_{EH} invariant under gauge transformations

$$g_{\mu\nu} \rightarrow g_{\mu\nu} + \nabla_{\mu}\zeta_{\nu} + \nabla_{\nu}\zeta_{\mu} + \nabla_{\mu}\zeta_{\lambda}\nabla_{\nu}\zeta^{\lambda}$$

- Introduce new fields $\bar{g}_{\mu\nu}$, ξ_{μ} and ϕ :

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + \nabla_{\mu}(\xi_{\nu} + \nabla_{\nu}\phi) + \nabla_{\nu}(\xi_{\mu} + \nabla_{\mu}\phi) \\ + \nabla_{\mu}(\xi_{\lambda} + \nabla_{\lambda}\phi) \cdot \nabla_{\nu}(\xi^{\lambda} + \nabla^{\lambda}\phi)$$

- Gauge invariances:

$$\bar{g}_{\mu\nu} \rightarrow \bar{g}_{\mu\nu} + \nabla_{\mu}\zeta_{\nu} + \dots, \quad \xi^{\mu} \rightarrow \xi^{\mu} - \zeta^{\mu}$$

and

$$\xi^{\mu} \rightarrow \xi^{\mu} + \nabla^{\mu}\psi, \quad \phi \rightarrow \phi - \psi$$

$$10 + 4 + 1 = 15 \text{ fields}$$

$$4 + 1 = 5 \text{ gauge invariances}$$

$15 - 2 \times 5 = 5$ propagating degrees of freedom,
right number for massive spin-2 field.

True if action for $\bar{g}_{\mu\nu}$, ξ^μ and ϕ is second order in derivatives

- Fierz–Pauli theory in Minkowski background: OK (only with FP mass term)
- Fierz–Pauli theory in curved background $g_{\mu\nu}^{(c)}$: **fourth order action for ϕ**

$$(g_{\mu\nu} - \eta_{\mu\nu})^2 \implies (\nabla_\mu \nabla_\lambda \phi \cdot \nabla_\nu \nabla^\lambda \phi) \cdot (g_{\mu\nu}^{(c)} - \eta_{\mu\nu})$$

6th, Boulware–Deser mode appears. It is a ghost.

NB: No matter what is the background (Einstein or not; small curvature or not)

Back to our model

Einstein backgrounds studied so far.

Massive vector field

$$u_{ij} = F_{(1)ij} - \frac{1}{6}\eta_{ij}F_{(1)}$$

$F_{(1)ij}$: perturbation of F_{ij} .

Linearized field equation ($\lambda = 0$ for brevity)

$$\begin{aligned} \nabla^2 u_{ij} - \nabla^k \nabla_i u_{kj} - \nabla^k \nabla_j u_{ki} + \nabla_i \nabla_j u + \eta_{ij} \left(\nabla^k \nabla^l u_{kl} - \nabla^2 u \right) \\ - m^2 (u_{ij} - \eta_{ij} u) + \text{const} \cdot W_{ilkj} u^{lk} = 0 \end{aligned}$$

W_{ilkj} : Weyl tensor of background

- First line: equation for massless tensor field in curved space-time;

invariance $u_{ij} \rightarrow u_{ij} + \nabla_i \zeta_j + \nabla_j \zeta_i$

No non-linear term

Stückelberg analysis

$$u_{ij} = \bar{u}_{ij} + \nabla_i \xi_j + \nabla_j \xi_i + \nabla_i \nabla_j \phi$$

- Mass term of Fierz–Pauli structure;
higher order terms in the action for ϕ cancel out
- Weyl term

$$S_W \propto \int d^4x \sqrt{-g} W_{iklj} u^{kl} u^{ij} \implies \int d^4x \sqrt{-g} W_{iklj} \nabla^k \nabla^l \phi \nabla^i \nabla^j \phi$$

Second order upon integration by parts:

$$S_W \propto \int d^4x \sqrt{-g} W_{iklj} W_{iklm} \nabla^j \phi \nabla^m \phi$$

No Boulware–Deser mode

Healthy theory at least in Einstein backgrounds
with small curvature.

A lot to be understood

- **Solutions.**
 - Is vDVZ problem cured by Vainshtein mechanism? Or one has to fine tune parameters to make massive spin-2 mode weakly interacting with matter?
 - Consistency with tests of GR
 - Do black holes have tensor hair?
- **Further consistency checks**
 - General backgrounds
 - Strong coupling scale in effective low energy theory
- **Cosmology**
- ...

