

---

**Bounds on new light  
particles from very small  
momentum transfer  
*np* elastic scattering data**

Yuri Kamyshkov, Jeffrey Tithof,

*University of Tennessee, TN 37996-1200, USA*

Mikhail Vysotsky

*ITEP, Moscow, 117218, Russia*

Phys.Rev.D78:114029,2008

4'th Sakharov Conference, FIAN, 18 May, 2009

---


---

NA-6 experiment, CERN SPS,

Results published in 1984

$E_n = 100 - 400$  GeV, gaseous hydrogen target

very small  $|t|$  (GeV)



---

$$\frac{d\sigma}{dt} = A \exp[bt] - 2 \left( \frac{\alpha k_n}{m_n} \right)^2 \frac{\pi}{t}, \quad (1)$$

where  $A = (79.78 \pm 0.26) \text{mb}/\text{GeV}^2$  and  $b = (11.63 \pm 0.08)$

$\text{GeV}^{-2}$  were determined from the fit to the data;

$k_n = -1.91$  is the neutron magnetic moment in nuclear magnetons;

factor “2” in the Schwinger term accounts for noncoherent sum of the scattering of neutron magnetic moment on proton and electron electric charges

$$\frac{d\sigma_i}{dt}(g, \mu)|_{\text{new}} = \frac{|A_i|^2}{16\pi s(s - 4m^2)} \quad , \quad (2)$$

where  $s = (p_n + p_p)^2$  is the invariant energy square and  $m$  is the nucleon mass.

---


$$|A_S|^2 = \frac{g_S^4}{(t - \mu^2)^2} (4m^2 - t)^2 \quad , \quad (3)$$

$$|A_P|^2 = \frac{g_P^4 t^2}{(t - \mu^2)^2} \quad , \quad (4)$$

$$|A_V|^2 = \frac{4g_V^4}{(t - \mu^2)^2} \left[ s^2 - 4m^2 s + 4m^4 + st + \frac{1}{2} t^2 \right] \quad , \quad (5)$$

$$|A_A|^2 = \frac{4g_A^4}{(t - \mu^2)^2} \left[ s^2 + 4m^2 s + 4m^4 + st + \frac{1}{2} t^2 + \frac{4m^4 t^2}{\mu^4} + \frac{8m^4 t}{\mu^2} \right] \quad , \quad (6)$$

where coupling constants  $g_i^2 \equiv g_p^i g_n^i$ .


$A \sim s^\alpha$ ;  $A_P \sim t$


---

Lack of fundamental theory for SI  $np$  scattering

amplitude does not prevent us from excluding light new particles as far as there are NO SI particles lighter than

pion,  $m_\pi = 140MeV$





90% C.L. bounds

---

Our bounds on the parameters  $g_V^2$  and  $g_A^2$  are rather strong; say for  $\mu = 10$  MeV,  $g_{V,A}^2 < 5 \cdot 10^{-3}$  at 90% C.L., which corresponds to

$$g_N^{V,A} < 0.071 \quad , \quad (7)$$

four times smaller than the QED coupling constant  $\sqrt{4\pi\alpha} \simeq 0.3$ . For scalar exchange, taking  $\mu = 10$  MeV, we get a much weaker bound,  $g_S^2 < 1.4$ .

---

It is quite natural to suppose that couplings of a new light particle with nucleons originate from its couplings with quarks. In this case  $A_V$  and  $A_A$  are modified. For vector exchange the induced magnetic moment interaction term should be added to the scattering amplitude. Since its numerator contains momentum transfer divided by  $m_N$  which in considered kinematics gives a factor much smaller than 1, we can safely neglect it.

The case of axial exchange is more delicate:

$$\begin{aligned}
\tilde{A}_A &= g_A^2 \bar{n} \gamma_\beta \gamma_5 n \left( g_{\alpha\beta} - \frac{k_\alpha k_\beta}{k^2 - m_\pi^2} \right) \frac{(g_{\alpha\mu} - \frac{k_\alpha k_\mu}{\mu^2})}{k^2 - \mu^2} * \\
&* \left( g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2 - m_\pi^2} \right) \bar{p} \gamma_\nu \gamma_5 p = \\
&= \frac{g_A^2}{k^2 - \mu^2} \left[ g_{\alpha\beta} - \frac{k_\alpha k_\beta}{\mu^2} \frac{(m_\pi^4 - 2\mu^2 m_\pi^2 + k^2 \mu^2)}{(k^2 - m_\pi^2)^2} \right] * \\
&* \bar{n} \gamma_\alpha \gamma_5 n \bar{p} \gamma_\beta \gamma_5 p \tag{8}
\end{aligned}$$

for massless pion the axial current is conserved, while term  $k_\alpha k_\beta / k^2 \longrightarrow m_N^2 / k^2$  in the amplitude leads to regular diff. crosssection at  $t \longrightarrow 0$ .

# literature 1

---

$$V(r) = -G_N \frac{m_1 m_2}{r} [1 + \alpha_G \exp(-r/\lambda)] \quad , \quad (9)$$

$$\alpha_G = \frac{g_{V,S}^2}{4\pi G_N m_p m_n} = 1.35 \cdot 10^{37} g_{V,S}^2 \quad , \quad \lg \alpha = \lg g_{V,S}^2 + 37.13 \quad (10)$$

# literature 2


---

low energy (KeV)  $n - {}^{208}\text{Pb}$  scattering

$$\frac{d\sigma}{d\Omega} = \frac{\sigma_0}{4\pi} [1 + \omega E \cos \theta] \quad , \quad (11)$$

$$|\Delta\omega| = \frac{16m_n^2}{\sqrt{\sigma_0/4\pi}} \frac{g_n^2}{4\pi} \frac{A}{\mu^4} \quad . \quad (12)$$

$g_{V,S}^2 < 4 \cdot 10^{-6}$  for 10MeV boson



# literature 3

---

Couplings of new light bosons with quarks are bounded by pion and kaon decays:

$$Br(\pi^0 \rightarrow \nu\nu) < 2.7 \cdot 10^{-7}$$

$$Br(\pi^0 \rightarrow \gamma\nu\nu) < 6 \cdot 10^{-4}$$

$$Br(K^+ \rightarrow \pi^+ + \nu\nu) < 2 \cdot 10^{-10}$$

Bounds on axial and scalar couplings are considerably stronger while the bound on vector coupling is comparable with those from  $np$  scattering

# literature 4

---

NP contribution to nuclear matter EOS; neutron stars  
Krivoruchenko, Simkovic, Faessler, 2009  
Stimulated by our paper (?)

the fact that the bounds are similar to ours is not surprising:  
 $\delta V \sim g^2 / \mu^2$ , so precision data on  $np$  - scattering compete  
with information from observations of neutron stars.

# Conclusions

---

- Electric neutrality of neutron allows to look for large distance NP effects in small angle neutron scattering data
- High energy scattering gives access to small values of vector and axial coupling constants
- Our bound on  $g_V$  is comparable with bounds from neutral pion and kaon decays
- Relation with ADS - neutrons